



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number : **0 293 063 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification :  
18.03.92 Bulletin 92/12

(51) Int. Cl.<sup>5</sup> : **C07D 473/30, C07D 233/90,  
A61K 31/52**

(21) Application number : **88300696.7**

(22) Date of filing : **27.01.88**

(54) **2-Phenylpurinone derivatives, processes for their preparation, and their use as well as intermediates.**

(30) Priority : **02.02.87 GB 8702300  
02.02.87 GB 8702301**

(43) Date of publication of application :  
30.11.88 Bulletin 88/48

(45) Publication of the grant of the patent :  
18.03.92 Bulletin 92/12

(84) Designated Contracting States :  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(56) References cited :  
**WO-A-86/00305  
DE-A- 1 545 574  
GB-A- 1 338 235  
E. SCHRÖDER et al.: "Arzneimittelchemie I",  
1976, G. Thieme Verlag, Stuttgart, DE, pages  
24-38, Chapter 7.1  
CHEMICAL ABSTRACTS, vol. 77, no. 15, 9th  
October 1972, page 438, abstract no. 101674t,  
Columbus, Ohio, US; & JP - A - 72 20 031  
CHEMICAL ABSTRACTS, vol. 69, no. 3, 15th  
July 1968, page 1001, abstract 10477z, Colum-  
bus, Ohio, US; & JP - A - 67 20 067**

(56) References cited :  
**CHEMICAL ABSTRACT, vol. 101, no. 7, 13th  
August 1984, page 613, abstract 54988a, Col-  
umbus, Ohio, US; Y.S. MOKRUSHIN et al.:  
"Synthesis and properties of analogs of  
5(4)-aminoimidazole-4(5)-carboxamide and  
purines. 14. Acylation of 5(4)-aminoimidazole  
derivatives"  
CHEMICAL ABSTRACTS, vol. 97, no. 3, 19th  
July 1982, page 750, abstract 24147g, Colum-  
bus, Ohio, US; A. PARKIN et al.: "Acyclic  
analogs of purine and imidazole nucleosides"  
PATENT ABSTRACTS OF JAPAN, vol. 6, no. 73  
(C-101)[951] 8th March 1982**

(73) Proprietor : **SMITH KLINE & FRENCH  
LABORATORIES LIMITED  
Mundells  
Welwyn Garden City Hertfordshire, AL7 1EY  
(GB)**

(72) Inventor : **Coates, William John  
45 Lodgefield  
Welwyn Garden City Hertfordshire AL7 1SD  
(GB)**

(74) Representative : **Waters, David Martin, Dr. et al  
Smith Kline & French Laboratories Ltd. Patent  
Department Mundells  
Welwyn Garden City Hertfordshire AL7 1EY  
(GB)**


EP 0 293 063 B1

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

## Description

The present invention relates to purinone derivatives and in particular to such compounds having a substituted phenyl group at the 2-position of the purinone ring. This invention further relates to processes for their preparation, intermediates in their preparation, their use as therapeutic agents and to pharmaceutical compositions containing them. The compounds of this invention are inhibitors of a calmodulin insensitive cyclic GMP phosphodiesterase and are of use in combatting such conditions wherein such inhibition is thought to be beneficial. The compounds of this invention are bronchodilators and are therefore of use in combatting chronic reversible obstructive lung diseases such as asthma and bronchitis. In addition the compounds of the present invention exhibit anti-allergic activity and are therefore of use in combatting allergic diseases such as allergic asthma, allergic rhinitis, urticaria and irritable bowel syndrome. Furthermore the compounds of this invention are vasodilators and are therefore of value in combatting angina, hypertension and congestive heart failure.

Accordingly the present invention provides compounds of the formula (1):



and pharmaceutically acceptable salts thereof, wherein

R¹ is C<sub>1-6</sub>alkyl or C<sub>2-6</sub>alkenyl, and

R² is hydrogen or hydroxy.

Suitably R¹ is C<sub>2-6</sub>alkyl for example ethyl, n-propyl, isopropyl, butyl, isobutyl or pentyl.

Suitably R¹ is C<sub>3-6</sub>alkenyl for example allyl, butenyl or pentenyl.

Preferably R¹ is n-propyl.

Suitably R² is hydrogen. Suitably R² is hydroxy.

Particular compounds of this invention are:

2-(2-propoxyphenyl)-6-purinone,

2-(2-ethoxyphenyl)-6-purinone,

2-(2-butoxyphenyl)-6-purinone,

2-(2-isobutoxyphenyl)-6-purinone,

2-(2-propoxyphenyl)purine-6,8-dione,

2-(2-methoxyphenyl)purine-6,8-dione,

2-(2-ethoxyphenyl)purine-6,8-dione,


2-(2-butoxyphenyl)purine-6,8-dione,

2-(2-isobutoxyphenyl)purine-6,8-dione, and

2-(2-allyloxyphenyl)purine-6,8-dione

and pharmaceutically acceptable salts thereof.

This invention covers all tautomeric forms of compounds of formula (1). For example the compound of the formula (1) wherein R² is hydroxy can exist in a tautomeric keto form:



Compounds of the formula (1) wherein R² is hydrogen may form pharmaceutically acceptable salts with acids such as hydrochloric, hydrobromic, sulphuric, phosphoric, acetic, citric, maleic, lactic, ascorbic, fumaric,

oxalic, methanesulphonic and ethanesulphonic acids.

Compounds of the formula (1) may form pharmaceutically acceptable salts with metal ions, such as alkali metals for example sodium and potassium, or with an ammonium ion.

In order to use a compound of the formula (1) or a pharmaceutically acceptable salt thereof for the treatment of humans and other mammals it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

Compounds of formula (1) and their pharmaceutically acceptable salts may be administered in standard manner for the treatment of the indicated diseases, for example orally, parenterally, trans-dermally, rectally, via inhalation or via buccal administration.

Compounds of formula (1) and their pharmaceutically acceptable salts which are active when given orally or via buccal administration can be formulated as liquids, syrups, tablets, capsules and lozenges. An oral liquid formulation will generally consist of a suspension or solution of the compound or salt in a liquid carrier for example, ethanol, glycerine or water with a flavouring or colouring agent. Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include magnesium stearate, starch, celluloses, lactose and sucrose. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example using the aforementioned carriers in a hard gelatin capsule shell. Where the composition is in the form of a soft gelatin shell capsule any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous gums, celluloses, silicates or oils and are incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of the compound or salt in a sterile aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil, for example polyethylene glycol, polyvinyl-pyrrolidone, lecithin, arachis oil, or sesame oil.

A typical suppository formulation comprises a compound of formula (1) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent, for example polymeric glycols, gelatins, cocoa-butter or other low melting vegetable waxes or fats.

Typical transdermal formulations comprise a conventional aqueous or non-aqueous vehicle, for example a cream, ointment, lotion or paste or are in the form of a medicated plaster, patch or membrane.

Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be administered in the form of an aerosol using a conventional propellant such as dichlorodifluoromethane or trichlorofluoromethane, are in the form of a powder for insufflation.

Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer to himself a single dose.

Each dosage unit for oral administration contains suitably from 0.001 mg/Kg to 3 mg/Kg, and preferably from 0.005 mg/Kg to 1.5 mg/Kg, and each dosage unit for parenteral administration contains suitably from 0.001 mg/Kg to 1 mg/Kg, of a compound of formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base.


The daily dosage regimen for oral administration is suitably about 0.001 mg/Kg to 12 mg/Kg, of a compound of formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base. The daily dosage regimen for parenteral administration is suitably about 0.001 mg/Kg to 4 mg/Kg, for example about 0.005 mg/Kg to 1 mg/Kg, of a compound of the formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base. The active ingredient may be administered as required, for example from 1 to 8 times a day or by infusion. The compositions of the invention are bronchodilators and are useful in chronic reversible obstructive lung disease for example asthma and bronchitis. The compositions of the present invention have vasodilator activity and are of use in the treatment of angina, hypertension and congestive heart failure. Such conditions can be treated by administration orally, topically, rectally, parenterally or by inhalation. For administration by inhalation dosages are controlled by a valve, are administered as required and for an adult are conveniently in the range 0.1-5.0 mg of a compound of the formula (1) or a pharmaceutically acceptable salt thereof.

The compounds of this invention may be co-administered with other pharmaceutically active compounds, for example in combination, concurrently or sequentially. Conveniently the compounds of this invention and the other active compound or compounds are formulated in a pharmaceutical composition. Examples of compounds which may be included in pharmaceutical compositions with the compounds of the formula (1) are bronchodilators such as sympathomimetic amines for example isoprenaline, isoetharine, salbutamol, phenylephrine and ephedrine or xanthine derivatives for example theophylline and aminophylline, anti-allergic agents for example disodium cromoglycate, histamine H<sub>1</sub>-antagonists, vasodilators for example hydralazine, angiotensin converting enzyme inhibitors for example captopril, anti-anginal agents for example isosorbide nitrate, glyceryl trinitrate and pentaerythritol tetranitrate, anti-arrhythmic agents for example quinidine, procainamide and lignocaine, calcium antagonists for example verapamil and nifedipine, diuretics such as thiazides and related compounds for example bendroflumazide, chlorothiazide, chlorothalidone, hydrochlorothiazide, and other diuretics

for example frusemide and triamterene, and sedatives for example nitrazepam, flurazepam and diazepam.

In another aspect the present invention provides a process for the preparation of a compound of the formula (1) or a pharmaceutically acceptable salt thereof, which process comprises :


a) for compounds wherein  $R^2$  is hydrogen, reacting a compound of the formula (2) :



wherein  $R^1$  is as hereinbefore defined, with a formylating agent;


b) for compounds wherein  $R^2$  is hydroxy, reacting a compound of the formula (2) as hereinbefore defined with a carbonylating agent;

c) for compounds wherein  $R^2$  is hydrogen, reacting a compound of the formula (3) :



wherein  $R^1$  is as hereinbefore defined with 4-amino-5-imidazolecarboxamide,

d) for compounds wherein  $R^2$  is hydrogen, cyclising a compound of the formula (4) :



wherein  $R^1$  is as hereinbefore defined;

and thereafter optionally forming a pharmaceutically acceptable salt.

The reaction between a compound of the formula (2) and a formylating agent is conveniently performed in the absence of a solvent or in the presence of a suitable solvent such as a  $C_{1-4}$  alcohol, pyridine or N-methylpyrrolidone, at ambient or elevated temperature, for example 50-250°C, preferably 100-200°C. Examples of formylating agents include formic acid,  $C_{1-4}$  alkyl formate, formamide,  $C_{1-4}$  alkyl formamide, formamidine,  $C_{1-4}$  alkyl formamidine or tri( $C_{1-4}$ ) alkyl orthoformate. Suitably a compound of the formula (2) is reacted with formamidine acetate in the presence of sodium acetate. Preferably the compound of the formula (2) is used in the form of an acid addition salt, for example the sulphate, and is reacted with an excess of a formylating agent, for example formamide.


The reaction between a compound of the formula (2) and a carbonylating agent is conveniently performed in the absence of a solvent or in a suitable solvent such as a halohydrocarbon, pyridine or toluene, at ambient or elevated temperature, for example 50-250°C. Suitable carbonylating agents include urea, di( $C_{1-4}$ ) alkyl carbonate,  $C_{1-4}$  alkyl chloroformate, phosgene, trichloromethyl chloroformate or carbonyldiimidazole.

Suitably a compound of the formula (3) is reacted with an acid addition salt of 4-amino-5-imidazolecarboxamide, for example the hydrochloride, in the absence of a solvent or in a suitable solvent such as a  $C_{1-4}$  alcohol, pyridine or N-methylpyrrolidone at an elevated temperature, for example 50-250°C.

Suitably a compound of the formula (4) is cyclised by heating at an elevated temperature, for example 50-150°C, in the presence of an acid or a base in a suitable solvent such as aqueous  $C_{1-4}$  alcohols, water, toluene,

a halohydrocarbon or acetonitrile.

A compound of the formula (4) can be prepared by reaction of 4-amino-5-imidazolecarboxamide with a compound of the formula (5) :



wherein L¹ is halo and R¹ is as hereinbefore defined.

Suitably L¹ is chloro or bromo. Suitably a compound of the formula (5) is reacted with 4-amino-5-imidazolecarboxamide at ambient or elevated temperature e.g. 50-100°C in a suitable solvent such as toluene, acetonitrile or a halohydrocarbon e.g. chloroform or dichloromethane, optionally in the presence of a base such as pyridine or triethylamine, to form a compound of the formula (4) which may be cyclised in situ to form a compound of the formula (1) wherein R² is hydrogen or may be isolated and thereafter cyclised as hereinbefore described.

Compounds of the formulae (2) and (3) are known or preparable in conventional manner from GB Patent No 1338235.

Pharmaceutically acceptable acid addition salts of the compounds of the formula (1) wherein R² is hydrogen may be prepared from the corresponding base of the compounds of the formula (1) in conventional manner. For example the base may be reacted with an acid in a C<sub>1-4</sub>alkanol, or an ion-exchange resin may be used. The salts of the compounds of the formula (1) may be interconverted using ion-exchange resins. Non-pharmaceutically acceptable salts are therefore of use as they can be converted to pharmaceutically acceptable salts.

Pharmaceutically acceptable base addition salts of the compounds of the formula (1) may be prepared by standard methods, for example by reacting a solution of the compound of the formula (1) with a solution of the base.

The following biological test methods, data and Examples serve to illustrate this invention.

#### Bronchodilatation - In vivo

Male guinea-pigs of the Dunkin Hartley strain (500 - 600g) were anaesthetised with Sagatal (pentobarbital sodium) (60 mg/kg). Airway resistance was measured using a modification of the classical Konzett-Rossier technique (Versuchsanordnung zu Untersuchungen an der Bronchialmuskulatur. Naunyn-Schmiedeberg's Arch. Exp. Path. Pharmac., vol 195: pp 71-74, (1940)). U46619 (9,11-methanoepoxy-PGH<sub>2</sub>) was infused i.v. at a rate of 2.5 nmol/min, this produced a steady state of bronchoconstriction (approximately 120% increase from basal airway resistance). The compound under test was administered by i.v. bolus injection, and the subsequent peak inhibition of bronchoconstriction recorded.

The dose of compound required to reduce the U46619-induced bronchoconstriction by 50% is given as the BD<sub>50</sub>. These results demonstrate in vivo anti-bronchoconstrictor activity.

COMPOUND OF EXAMPLE	BD <sub>50</sub> (μmol/kg)
1	3.6
2	6.5
3	3.8
4	5.3
5	1.0
7	3.6
8	1.5
9	4.5

#### 20 Vasodilatation - in vivo

Male Wistar rats (300 g) were anaesthetised with a sodium 5-ethyl-5-(1-methylpropyl)-2-thiobarbiturate/sodium pentobarbitone mixture i.p. (62.5 and 22.5 mg/kg respectively). The trachea was cannulated and the rats breathed spontaneously air enriched with O<sub>2</sub> (5 ml/min). Blood pressure was recorded from a carotid artery and a jugular vein was cannulated for the administration of compounds. The temperature of the animal was maintained at 37°C by the use of an electric blanket. The abdominal aorta was separated from the inferior vena cava, distal to the renal arteries and was cannulated centrally to supply the perfusion pump with blood and distally for the perfusion of the hind quarters at constant pressure. The perfusion circuit was primed with 5% bovine serum albumin dissolved in 0.9% sodium chloride solution, pH adjusted to 7.4. Initially the pump rate was set between 10 and 15 ml/min to match the hind quarter perfusion pressure to that of the systemic circulation. Once set the pressure remained unaltered for the rest of the experiment. A change in the speed of the pump (equivalent to hindquarter blood flow) was used to assess the changes in hindquarter vascular resistance.

All compounds were administered as a bolus i.v. and from the dose response curves the dose required to produce a 50% increase in hindquarter blood flow (EDHQ<sub>50</sub>) was determined in μmoles/kg. The following results were obtained :

COMPOUND OF EXAMPLE	EDHQ <sub>50</sub> (μmol/kg)
1	10.8
5	3.7

#### 45 Anti-allergic activity - in vitro

Male Hartley strain guinea pigs (weighing 250-300 g) were actively sensitized to ovalbumin (OA) by a modified Herxheimer method (J. Physiol. 117:251-258, 1952). The animals were injected intramuscularly with 0.7 ml of a 5.0% OA solution prepared in isotonic saline. Four weeks after sensitization tracheas were removed and cut into spiral strips. Each trachea was cut in half, placed in 10-ml water-jacketed tissue baths and attached with biological tissue clips via silk suture to force displacement transducers for recording isometric tension. One half of each trachea served as a control for the corresponding drug-treated tissue. The tracheas were bathed in modified Krebs-Henseleit solution of the following composition (mM): NaCl, 118; KCl, 4.7; MgSO<sub>4</sub>, 1.2; CaCl<sub>2</sub>, 2.5; NaHCO<sub>3</sub>, 25; KH<sub>2</sub>PO<sub>4</sub>, 1.2; and glucose, 10. The physiologic buffer was maintained at 37°C and continually aerated with 95% O<sub>2</sub>/5% CO<sub>2</sub>. The tissue preparations were placed under 2 g passive tension and equilibrated

for 60 minutes, during which time they were washed every 15 minutes with fresh buffer. Each tissue was pretreated for 45 minutes with meclofenamic acid ( $1 \mu\text{M}$ ).

Tissues were pretreated with the compound under test ( $100 \mu\text{M}$ ), or vehicle ( $0.4 \text{ mM NaOH}$ ) for 30 minutes prior to the addition of OA. OA ( $0.1 \mu\text{g/ml}$ ) was added to all tissues and the contraction was monitored for 15 minutes. Previous experiments indicated that this concentration of OA produced the maximum antigen-induced contraction. At the conclusion of the experiment, a maximally-effective concentration of carbachol ( $10 \mu\text{M}$ ) was added to each tissue and OA-induced responses were expressed as a percentage of this reference contraction. Since the compounds under test reduced basal tone, the absolute response (i.e., g tension) to carbachol was calculated as the difference between the tension after the addition of the test compounds and the maximum tension developed in the presence of carbachol. To evaluate effects of the test compounds, the degree of contraction was calculated 2 and 12 minutes after the addition of OA. OA-induced contraction of drug-treated tissues was expressed as a percentage of the response of vehicle-treated, paired controls.

When administered prior to OA challenge, both the compound of Example 1 ( $100 \mu\text{M}$ ) and the compound of Example 5 ( $100 \mu\text{M}$ ) significantly reduced basal tone. As a percentage of the maximum response to carbachol, the compound of Example 1 reduced basal tone by  $17 \pm 3\%$  and the compound of Example 5 by  $24 \pm 7\%$ . Administration of OA to vehicle-treated control tissues produced contractions which reached a maximum ( $67 \pm 2\%$  of the maximum response to carbachol;  $N=4$ ) within 3 minutes and were sustained over the 15-minute observation period. Two minutes after the addition of OA, the response of tissues treated for 30 minutes with the compound of Example 1 ( $100 \mu\text{M}$ ) was  $78 \pm 12\%$  (N.S.,  $P > 0.05$ ;  $N=4$ ) of the control and the contractile response of those treated with the compound of Example 5 ( $100 \mu\text{M}$ ) was  $79 \pm 3\%$  ( $P < 0.05$ ;  $N=4$ ) of the control. Twelve minutes after OA challenge, contractile force had declined ( $P < 0.05$ ) in tissues treated with either the compound of Example 1 or the compound of Example 5 to  $46 \pm 9\%$  or  $60 \pm 4\%$ , respectively, of the control values.

Results are expressed as a mean  $\pm$  S.E. of 4 experiments. Statistical significance of differences between the means of control and drug-treated groups was determined using a paired Student's *t* test.

These results indicate that compounds of Examples 1 and 5 inhibit antigen-induced contraction of the guinea-pig isolated trachea.

#### Anti-allergic activity - in vivo

Adult male albino Hartley strain guinea pigs weighing 400-600 g were actively sensitized to OA according to the procedure hereinbefore described. To measure pulmonary mechanics, a polyethylene catheter was inserted into the trachea of anesthetized (sodium pentobarbital,  $35 \text{ mg/kg}$ , i.p.) animals and airflow was measured via a heated pneumotachograph in combination with a differential pressure transducer connected to the pneumotachograph. Airflow and transpulmonary pressure signals were fed into an on-line pulmonary mechanics computer which integrated the flow signal to obtain tidal volume. Total pulmonary resistance ( $R_L$ ) and dynamic lung compliance ( $C_{DYN}$ ) were calculated at isovolumetric points according to the method of Amdur and Mead (Am. J. Physiol. 192:364-368, 1958).

A compound under test was dissolved in  $25 \text{ mM NaHCO}_3$  plus one drop of polyethylene glycol 400 to a concentration of  $100 \text{ mM}$ . Based on the weight of the animal, the appropriate amount of solution was removed and the volume brought up to  $1 \text{ ml}$  with  $25 \text{ mM NaHCO}_3$ . For the intraduodenal administration of test compounds, a small midline incision was made in the abdomen, the duodenum was exposed, and a butterfly needle attached to plastic tubing was inserted. All solutions were adjusted to the same volume ( $1 \text{ ml}$ ) and administered through the butterfly needle and followed with  $0.5 \text{ ml}$  of distilled water. The needle was then removed and the abdominal incision was stapled to prevent heat loss.

Aerosols of OA were generated to deliver approximately  $0.375 \mu\text{l/breath}$ . Pressure, flow rate and sensitivity were set such that a maximum pleural pressure of  $20 \text{ cm water}$  was attained during inhalation. One hour after the administration of the test compounds, guinea pigs were challenged with 5 inhalations of OA ( $0.5 \text{ mg/ml}$ ) via the tracheal cannula and changes in  $C_{DYN}$  and  $R_L$  were monitored for 10 minutes. Baseline values for  $C_{DYN}$  and  $R_L$  were  $0.4\text{-}0.6 \text{ ml/cm water}$  and  $0.1\text{-}0.2 \text{ cm water/ml/sec}$ , respectively.

Administration of OA to vehicle-treated, anesthetized guinea pigs produced bronchoconstriction which developed slowly over a 5-minute period. Five minutes after OA challenge,  $R_L$  had increased  $162 \pm 40\%$  and  $C_{DYN}$  had decreased  $51 \pm 26\%$  ( $N=5$ ). These changes in pulmonary function were stable over the remaining 5 minutes of the observation period. Intraduodenal administration of  $100 \mu\text{mol/kg}$  of the compound of Example 1 ( $N=4$ ) 60 minutes before OA challenge significantly reduced antigen-induced bronchoconstriction: 5 minutes after challenge,  $C_{DYN}$  was unchanged from baseline ( $P < 0.01$  vs. vehicle-treated controls) and the increase in  $R_L$  was limited to  $51 \pm 26\%$  ( $P < 0.05$  vs. vehicle-treated controls).

Results are expressed as a mean  $\pm$  S.E. of 4 experiments. Statistical significance of differences between

the means of control and drug-treated groups was determined using a unpaired Student's t test.

### Inhibition of Phosphodiesterase (PDE)

5 Eight pig hearts/lungs were collected from the abattoir and kept on ice until the pulmonary arteries or aortas could be dissected and excess fat removed. 120 g of tissue was dissected. Unless otherwise stated all procedures were done at 4° C. Following dissection arteries were flash frozen in liquid nitrogen and stored at -70° C until required. On the day of homogenisation, tissue was cooled with liquid nitrogen and broken into small pieces by striking with a hammer. The tissue was then homogenised in 15 mM BIS-TRIS, 1 µg/ml leupeptin and antipain, 2 µg/ml pepstatin A, 5 µM benzamidine, 2 mM EDTA (ethylenediaminetetraacetic acid) and 2 mM dithiothreitol, pH 6.5. Phenylmethanesulphonylfluoride (PMSF) was added to a final concentration of 50 µM just prior to homogenisation. The homogenate was then centrifuged for 20 minutes at 30,000 g. Supernatant was filtered firstly through glass wool and then a 0.45 µm filter. The resultant filtrate was then applied to a 60 ml DEAE-Sepharose<sup>®</sup> CL-6B (Diethylaminoethyl Cellulose with a bead size of 45-165 microns) (Pharmacia) column pre-equilibrated in homogenisation buffer. The column was washed with 150 mls of homogenisation buffer and PDE activities eluted with 150 ml of homogenisation buffer containing 100 mM sodium acetate. Six 25 ml fractions were collected, the flow rate throughout was 80 ml/hr.

Fractions 2 and 3 were pooled and BIS-TRIS, MgCl<sub>2</sub>, CaCl<sub>2</sub> added to final concentrations of 50 mM, 7 mM and 5 mM respectively. PMSF, leupeptin, antipain and pepstatin A were also added at the concentrations described above for these components. The pH of this pooled sample was corrected to 6.9.

The sample was applied, at a flow rate of 3 ml/hr, to a tandem column set up. The first column in the tandem was a 5 ml calmodulin-agarose column (Sigma) and the second a 1.5 ml cibacron blue 3GA-agarose column (Sigma). Both columns were pre-equilibrated with 50 mM BIS-TRIS, 5 mM MgCl<sub>2</sub>, 5 mM CaCl<sub>2</sub>, 5 mM benzamidine, 2 mM dithiothreitol pH 6.9. Following application of the sample the columns were washed with 20 ml of equilibration buffer.

The cibacron blue-agarose column was then disconnected and washed with a variety of buffers as indicated below (Buffer A = 50 mM BIS-TRIS, 5mM benzamidine, 2 mM dithiothreitol pH 7.0).

20 ml Buffer A + 2 M NaCl, 5 mM MgCl<sub>2</sub>

20 ml Buffer A + 2 M NaCl

30 20 ml Buffer A + 2 M NaCl, 10 mM EDTA

PDE activity bound to the column was eluted with eight 2.5 ml aliquots of Buffer A containing 2 M NaCl, 10 mM EDTA and 10 mM cGMP. Fractions found to contain PDE activity were pooled and concentrated to approximately 1/3 the original volume by dialysis for 2 hr against Buffer A:glycerol (50:50) containing 15 mM 2-mercaptoethanol, 2 mM EDTA. After dialysis the concentration of cGMP in the fraction was reduced by treatment using a PD-10 column (Pharmacia) using dialysis buffer.

The fraction produced was stored at -20°C and for experiments diluted 50-fold and used as the source of PDE activity for the assay of compounds.

### Properties of PDE activity

40 The PDE activity isolated by the procedure described above demonstrated considerable selectivity for cGMP as a substrate showing only poor rates of hydrolysis of cAMP. The PDE activity of this preparation could not be increased by the presence of Ca<sup>2+</sup> (10 mM) and calmodulin (25 µg) in incubations and the hydrolysis of cAMP was not increased by the addition of 10 µM cGMP to incubations. The kinetic characteristics of this enzyme are summarised below.

$K_m$  cGMP = 1.3 µM

$K_m$  cAMP >120 µM

Ratio of cGMP/cAMP hydrolysis at 1 µM substrate = 30.

### Phosphodiesterase Assay

The assay was as described by Davis & Daly (1979) J. Cyclic Nucleotide Res., 5, 65-74, but with some modifications. The standard reaction mixture contained, in a final volume of 100 µl,

50 µM 5'-GMP (including 4000 dpm <sup>14</sup>C-GMP)

55 1 µM 3', 5'-cGMP (including 0.1 µCi <sup>3</sup>H-cGMP)

10 µM enzyme preparation

10 µM inhibitor dilution

and buffered with 50 mM TRIS/5 mM MgCl<sub>2</sub> pH 7.5. The reaction was initiated with enzyme, and was carried



out at 37°C for 5-10 minutes. The reaction was terminated by placing tubes in a boiling water bath for 2 minutes. 500  $\mu$ M of 0.1 M HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid) buffer pH 8.5 containing 0.1 M NaCl was then added to each assay tube and the contents applied to Affigel 601 (Bio-Rad) boronate affinity chromatography media (1.2 ml bed volume) which had previously been equilibrated with 10 ml of HEPES/NaCl buffer. Unreacted  $^3\text{H}$ -cGMP was washed from the column with  $10 \times 1$  ml HEPES/NaCl buffer. Labelled 5'-AMP was eluted into a scintillation vial with 6 ml 0.25 M acetic acid and counted in a scintillation counter using 10 ml Instagel (Packard). Recoveries were between 50%-75% as measured by recovery of  $^{14}\text{C}$ -GMP. Assays were performed in duplicate and values corrected for blanks of <2%.

#### Calculation of $\text{IC}_{50}$ values

$\text{IC}_{50}$  values (the concentration of inhibitor required for 50% inhibition of activity) were obtained by incubation of the enzymes at 1  $\mu$ M cyclic GMP and a range of inhibitor concentrations.

COMPOUND OF EXAMPLE	$\text{IC}_{50}$ ( $\mu\text{M}$ )
1	0.96
2	4.49
3	5.04
4	1.74
5	1.25
7	5.26
8	6.79
9	3.64

#### Example 1

##### 2-(2-Propoxyphenyl)-6-purinone

A stirred mixture of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one sulphate (1.5 g) (prepared by the addition of concentrated sulphuric acid to an ethanolic solution of the free base) and formamide (15 ml) was heated in an oil bath (temp. 190°-200°C) for 70 minutes. When cool the mixture was filtered and the collected solid was washed with ethanol to give a crude product (1.1 g), m.p. 254-259°C, which was recrystallised from ethanol to give the title compound, 0.72 g, m.p. 263-265°C.

#### Example 2

##### 2-(2-Ethoxyphenyl)-6-purinone

In a similar manner to Example 1 reaction of 4,5-diamino-2-(2-ethoxyphenyl)pyrimidin-6-one sulphate (1.5 g) with formamide (15 ml) afforded the title compound, 0.36 g, m.p. 276-277°C, (recrystallised from ethanol).

#### Example 3

##### 2-(2-Butoxyphenyl)-6-purinone

In a similar manner to Example 1 reaction of 4,5-diamino-2-(2-butoxyphenyl)pyrimidin-6-one sulphate (1.5 g) with formamide (5 ml) afforded the title compound, 0.65 g, m.p. 247-248°C, (recrystallised from ethanol).

**Example 4****2-(2-Isobutoxyphenyl)-6-purinone**

5 In a similar manner to Example 1 reaction of 4,5-diamino-2-(2-isobutoxyphenyl)pyrimidin-6-one sulphate (1.4 g) with formamide (5 ml) afforded the title compound, 0.24 g, m.p. 272-273°C, (recrystallised from ethanol).

**Example 5****2-(2-Propoxyphenyl)purine-6,8-dione**

10 A mixture of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one (1.3 g), and urea (1.5 g) was heated in an oil bath (temp. 190°C) for 45 minutes. The resultant solid was digested with hot water, the mixture filtered and the solid washed with water to give a crude product, 1.36 g. Recrystallisation from dimethylformamide gave  
 15 the title compound (1.01 g), m.p. >350°C,  $\delta$ (DMSO- $d_6$ ), 1.01 (t, 3H); 1.88 (m, 2H); 4.09 (t, 2H); 7.10, 7.21, 7.52 and 7.76 (multiplets, 4H);  $\tau$  11.07, 11.55 and 11.95 (very broad singlets, 3H).

**Example 6****2-(2-Methoxyphenyl)purine-6,8-dione**

20 In a similar manner to Example 5 reaction of 4,5-diamino-2-(2-methoxyphenyl)pyrimidin-6-one (0.93 g) with urea (1.20 g) afforded the title compound, 0.28 g, m.p. 329-330°C, (recrystallised twice from dimethylformamide).

**Example 7****2-(2-Ethoxyphenyl)purine-6,8-dione**

30 A solution of 4,5-diamino-2-(2-ethoxyphenyl)pyrimidin-6-one sulphate (2.0 g) in water (50 ml) was neutralized with ammonium hydroxide and extracted with chloroform. The organic extract was evaporated under reduced pressure to dryness and the residual free base was treated in a similar manner to Example 5 with urea (1.74 g) to afford after recrystallisation from dimethylformamide the title compound, 0.66 g, m.p. 349-351°C.

**Example 8****2-(2-Butoxyphenyl)purine-6,8-dione**

40 In a similar manner to Example 5 reaction of 4,5-diamino-2-(2-butoxyphenyl)pyrimidin-6-one (0.96 g) with urea (1.05 g) afforded the title compound, 0.26 g, m.p. 324-326°C, (recrystallised from dimethylformamide).

**Example 9****2-(2-Isobutoxyphenyl)purine-6,8-dione**

45 Carbonyldiimidazole (1.01 g) was added to 4,5-diamino-2-(2-isobutoxyphenyl)pyrimidin-6-one (1.50 g) in toluene (100 ml) and the resulting mixture was heated under reflux for one hour yielding a brown solid which was collected. This solid was washed with water and recrystallised from dimethylformamide to yield the title compound, 0.22 g, m.p. >360°C.

**Example 10****2-(2-Allyloxyphenyl)purine-6,8-dione**

55 4,5-Diamino-2-(2-allyloxyphenyl)pyrimidin-6-one sulphate (0.8 g) was added to a stirred solution of carbonyldiimidazole (0.72 g) in dry pyridine (8 ml) under nitrogen. The resulting solution was stirred under nitrogen for 3 hours at ambient temperature and then the volume of the solution was reduced by evaporation under reduced pressure. The residual syrup was diluted with aqueous acetone (50%, 20 ml) and the resultant solid

was collected, washed with water and acetone and dried to give a crude product (0.48 g). This material together with another crude sample (0.1 g, prepared in a similar manner as hereinbefore described) was recrystallised from acetic acid (ca 25 ml) by the addition of hot water (5 ml) to afford, after washing with acetic acid, water and ethanol and drying, the title compound, 0.4 g, m.p. 340-345°C dec.,  $\delta$ (DMSO- $d_6$ ), 4.68 (m, 2H); 5.2-5.4 (m, 2H); 5.9-6.2 (m, 1H); 7.07-7.2, 7.4-7.55, 7.6-7.7 (m's, 4H); 10.9, 11.45 and 12.1 (broad singlets, 3H).

#### Example 11

##### 2-(2-Propoxyphenyl)-6-purinone

A mixture of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one (0.3 g), formamidine acetate (0.18 g) and anhydrous sodium acetate (0.1 g) was heated in an oil bath at 155°-165°C for 2½ hours. The mixture melted and then resolidified. Ethanol (1 ml) was added and the title compound was collected by filtration, 0.31 g, m.p. 256-258°C.

#### Example 12

##### 2-(2-Propoxyphenyl)purine-6,8-dione

In a similar manner to Example 9 reaction of carbonyldiimidazole (4.98 g) with 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one (7.00 g) in toluene (350 ml) afforded the title compound, 3.76 g, m.p. >340°C (recrystallised from dimethylformamide).

#### Example 13

Pharmaceutical compositions for oral administration are prepared by combining the following :

	% w/w		
2-(2-Propoxyphenyl)-6-purinone	0.5	3.0	7.14
2% w/w Soya lecithin in soya bean oil	90.45	88.2	84.41
Hydrogenated vegetable shortening and beeswax	9.05	8.8	8.45

The formulations are then filled into individual soft gelatin capsules.


#### Example 14

A pharmaceutical composition for parenteral administration is prepared by dissolving the title compound of Example 1 (0.02 g) in polyethylene glycol 300 (25 ml) with heating. This solution is then diluted with water for injections Ph. Eur. (to 100 ml). The solution is then sterilised by filtration through a 0.22 micron membrane filter and sealed in sterile containers.

## Claims

Claims for the following Contracting State : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. A compound of the formula (1) :



or a pharmaceutically acceptable salt thereof, wherein  
R¹ is C₁-₆alkyl or C₂-₆alkenyl, and  
R² is hydrogen or hydroxy.


2. A compound according to claim 1 wherein R² is hydrogen;  
3. A compound according to claim 1 wherein R² is hydroxy.  
4. A compound according to any one of claims 1 to 3 wherein R¹ is C₂-₆alkyl.  
5. A compound according to any one of claims 1 to 3 wherein R¹ is C₃-₆alkenyl.  
6. A compound according to claim 1 which is :

2-(2-propoxyphenyl)-6-purinone,  
2-(2-ethoxyphenyl)-6-purinone,  
2-(2-butoxyphenyl)-6-purinone,  
2-(2-isobutoxyphenyl)-6-purinone,  
2-(2-propoxyphenyl)purine-6,8-dione,  
2-(2-methoxyphenyl)purine-6,8-dione,  
2-(2-ethoxyphenyl)purine-6,8-dione,  
2-(2-butoxyphenyl)purine-6,8-dione,  
2-(2-isobutoxyphenyl)purine-6,8-dione, or  
2-(2-allyloxyphenyl)purine-6,8-dione  
or a pharmaceutically acceptable salt thereof.

7. 2-(2-Propoxyphenyl)-6-purinone or a pharmaceutically acceptable salt thereof.  
8. 2-(2-Propoxyphenyl)purine-6,8-dione or a pharmaceutically acceptable salt thereof.  
9. A compound according to any one of claims 1 to 8 for use as a medicament.  
10. A compound according to any one of claims 1 to 8 for use as a bronchodilator.  
11. A compound according to any one of claims 1 to 8 for use as a vasodilator.  
12. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 8 and a pharmaceutically acceptable carrier.


13. A process for preparing a compound of the formula (1) or a pharmaceutically acceptable salt thereof as defined in claim 1 which comprises :

- a) for compounds wherein R² is hydrogen, reacting a compound of the formula (2) :




- wherein R¹ is as defined in claim 1, with a formylating agent;  
b) for compounds wherein R² is hydroxy, reacting a compound of the formula (2) as hereinbefore defined with a carbonylating agent;

c) for compounds wherein R<sup>2</sup> is hydrogen, reacting a compound of the formula (3) :




10 wherein R<sup>1</sup> is as hereinbefore defined with 4-amino-5-imidazolecarboxamide,  
d) for compounds wherein R<sup>2</sup> is hydrogen, cyclising a compound of the formula (4) :



wherein R<sup>1</sup> is as hereinbefore defined; and thereafter optionally forming a pharmaceutically acceptable salt.


14. A compound of the formula (4) :



wherein R<sup>1</sup> is C<sub>1-6</sub>alkyl or C<sub>2-6</sub>alkenyl.

35 **Claims for the following Contracting States : ES, GR**

1. A process for preparing a compound of the formula (1) :



50 or a pharmaceutically acceptable salt thereof, wherein


R<sup>1</sup> is C<sub>1-6</sub>alkyl or C<sub>2-6</sub>alkenyl, and

R<sup>2</sup> is hydrogen or hydroxy,


which process comprises :

a) for compounds wherein R<sup>2</sup> is hydrogen; reacting a compound of the formula (2) :


55



10 wherein R¹ is as hereinbefore defined, with a formylating agent;  
 b) for compounds wherein R² is hydroxy,  
 reacting a compound of the formula (2) as hereinbefore defined with a carbonylating agent;  
 c) for compounds wherein R² is hydrogen, reacting a compound of the formula (3) :



wherein R¹ is as hereinbefore defined with 4-amino-5-imidazolecarboxamide,  
 d) for compounds wherein R² is hydrogen, cyclising a compound of the formula (4) :



- wherein R¹ is as hereinbefore defined;
- 35 and thereafter optionally forming a pharmaceutically acceptable salt.
2. A process according to claim 1 for preparing a compound wherein R² is hydrogen.
  3. A process according to claim 1 for preparing a compound wherein R² is hydroxy.
  4. A process according to any one of claims 1 to 3 for preparing a compound wherein R¹ is C₂-₆alkyl.
  5. A process according to any one of claims 1 to 3 for preparing a compound wherein R¹ is C₃-₆alkenyl.
  - 40 6. A process according to claim 1 for preparing a compound which is :
    - 2-(2-propoxyphenyl)-6-purinone,
    - 2-(2-ethoxyphenyl)-6-purinone,
    - 2-(2-butoxyphenyl)-6-purinone,
    - 2-(2-isobutoxyphenyl)-6-purinone,
    - 45 2-(2-propoxyphenyl)purine-6,8-dione,
    - 2-(2-methoxyphenyl)purine-6,8-dione,
    - 2-(2-ethoxyphenyl)purine-6,8-dione,
    - 2-(2-butoxyphenyl)purine-6,8-dione,
    - 2-(2-isobutoxyphenyl)purine-6,8-dione, or
    - 50 2-(2-allyloxyphenyl)purine-6,8-dione
 or a pharmaceutically acceptable salt thereof.
  7. A process according to claim 1 wherein a formylating agent is selected from formic acid, C₁-₄alkyl formate, formamide, C₁-₄alkyl formamide, formamidine, C₁-₄alkyl formamidine or tri(C₁-₄)alkyl orthoformate.
  8. A process according to claim 1 wherein a carbonylating agent is selected from urea, di(C₁-₄)alkyl carbonate, C₁-₄alkyl chloroformate, phosgene, trichloromethyl chloroformate or carbonyldiimidazole.
  - 55 9. A process for preparing 2-(2-propoxyphenyl)-6-purinone or a pharmaceutically acceptable salt thereof which comprises reacting 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one or an acid addition salt thereof with

formamide or formamidine acetate and optionally forming a pharmaceutically acceptable salt thereof.


10. A process for preparing 2-(2-propoxyphenyl)purine-6,8-dione or a pharmaceutically acceptable salt thereof which comprises reacting 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one with urea or carbonyldiimidazole and optionally forming a pharmaceutically acceptable salt thereof.

11. A process for preparing a pharmaceutical composition which comprises bringing into association a compound of the formula (1) as defined in claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

## Patentansprüche

Patentansprüche für folgende Vertragsstaaten :AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Verbindung der Formel (1):



oder ein pharmazeutisch verträgliches Salz davon, in der  
R¹ einen C<sub>1-6</sub>-Alkyl- oder einen C<sub>2-6</sub>-Alkenylrest bedeutet, und  
R² ein Wasserstoffatom oder eine Hydroxylgruppe ist.

2. Verbindung gemäß Anspruch 1, wobei R² ein Wasserstoffatom bedeutet.

3. Verbindung gemäß Anspruch 1, wobei R² eine Hydroxylgruppe bedeutet.

4. Verbindung gemäß einem der Ansprüche 1 bis 3, wobei R¹ einen C<sub>2-6</sub>-Alkylrest bedeutet.

5. Verbindung gemäß einem der Ansprüche 1 bis 3, wobei R¹ einen C<sub>3-6</sub>-Alkenylrest bedeutet.

6. Verbindung gemäß Anspruch 1, die

2-(2-Propoxyphenyl)-6-purinon

2-(2-Ethoxyphenyl)-6-purinon

2-(2-Butoxyphenyl)-6-purinon

2-(2-Isobutoxyphenyl)-6-purinon

2-(2-Propoxyphenyl)purin-6,8-dion

2-(2-Methoxyphenyl)purin-6,8-dion

2-(2-Ethoxyphenyl)purin-6,8-dion

2-(2-Butoxyphenyl)purin-6,8-dion

2-(2-isobutoxyphenyl)purin-6,8-dion oder

2-(2-Allyloxyphenyl)purin-6,8-dion

oder ein pharmazeutisch verträgliches Salz davon bedeutet.

7. 2-(2-Propoxyphenyl)-6-purinon oder ein pharmazeutisch verträgliches Salz davon.

8. 2-(2-Propoxyphenyl)purin-6,8-dion oder ein pharmazeutisch verträgliches Salz davon.

9. Verbindung gemäß einem der Ansprüche 1 bis 8 zur Verwendung als Arzneimittel.


10. Verbindung gemäß einem der Ansprüche 1 bis 8 zur Verwendung als Bronchodilatator.

11. Verbindung gemäß einem der Ansprüche 1 bis 8 zur Verwendung als Vasodilatator.

12. Arzneimittel, umfassend eine Verbindung gemäß einem der Ansprüche 1 bis 8 und einen pharmazeutisch verträglichen Träger.


13. Verfahren zur Herstellung einer Verbindung der Formel (1) oder eines pharmazeutisch verträglichen Salzes davon, wie in Anspruch 1 definiert, umfassend:

a) für Verbindungen, in denen R² ein Wasserstoffatom bedeutet, die Umsetzung einer Verbindung der Formel (2):




(2)

in der R¹ wie in Anspruch 1 definiert ist, mit einem Formylierungsmittel,  
b) für Verbindungen, in denen R² eine Hydroxylgruppe bedeutet, die Umsetzung einer Verbindung der Formel (2), wie vorstehend definiert, mit einem Carbonylierungsmittel,  
c) für Verbindungen, in denen R² ein Wasserstoffatom bedeutet, die Umsetzung einer Verbindung der Formel (3):



(3)


in der R¹ wie vorstehend definiert ist, mit 4-Amino-5-imidazolcarboxamid,  
d) für Verbindungen, in denen R² ein Wasserstoffatom bedeutet, die Cyclisierung einer Verbindung der Formel (4):



(4)

in der R¹ wie vorstehend definiert ist, und danach gegebenenfalls die Bildung eines pharmazeutisch verträglichen Salzes.

14. Verbindung der Formel (4):




(4)

in der R¹ einen C₁-₆-Alkyl- oder einen C₂-₆-Alkenylrest bedeutet.

Patentansprüche für folgenden Vertragsstaaten : ES, GR

1. Verfahren zur Herstellung einer Verbindung der Formel (1):






(1)

oder eines pharmazeutisch verträglichen Salzes davon, in der R¹ einen C<sub>1-6</sub>-Alkyl- oder einen C<sub>2-6</sub>-Alkenylrest bedeutet, und R² ein Wasserstoffatom oder eine Hydroxylgruppe ist, welches Verfahren

a) für Verbindungen, in denen R² ein Wasserstoffatom bedeutet, die Umsetzung einer Verbindung der Formel (2):




(2)

in der R¹ wie vorstehend definiert ist, mit einem Formylierungsmittel,

b) für Verbindungen, in denen R² eine Hydroxylgruppe bedeutet, die Umsetzung einer Verbindung der Formel (2), wie vorstehend definiert, mit einem Carbonylierungsmittel,


c) für Verbindungen, in denen R² ein Wasserstoffatom bedeutet, die Umsetzung einer Verbindung der Formel (3):



(3)

in der R¹ wie vorstehend definiert ist, mit 4-Amino-5-imidazolcarboxamid,

d) für Verbindungen, in denen R² ein Wasserstoffatom bedeutet, die Cyclisierung einer Verbindung der Formel (4):



(4)

in der R¹ wie vorstehend definiert ist, und danach gegebenenfalls die Bildung eines pharmazeutisch verträglichen Salzes umfaßt.

2. Verfahren gemäß Anspruch 1 zur Herstellung einer Verbindung, in der R² ein Wasserstoffatom bedeutet.

3. Verfahren gemäß Anspruch 1 zur Herstellung einer Verbindung, in der R² eine Hydroxylgruppe bedeutet.

4. Verfahren gemäß einem der Ansprüche 1 bis 3 zur Herstellung einer Verbindung, in der R<sup>1</sup> einen C<sub>2-5</sub>-Alkylrest bedeutet.

5. Verfahren gemäß einem der Ansprüche 1 bis 3 zur Herstellung einer Verbindung, in der R<sup>1</sup> einen C<sub>3-6</sub>-Alkenylrest bedeutet.

5 6. Verfahren gemäß Anspruch 1, zur Herstellung einer Verbindung, die:

2-(2-Propoxyphenyl)-6-purinon

2-(2-Ethoxyphenyl)-6-purinon

2-(2-Butoxyphenyl)-6-purinon

2-(2-Isobutoxyphenyl)-6-purinon

10 2-(2-Propoxyphenyl)purin-6,8-dion

2-(2-Methoxyphenyl)purin-6,8-dion

2-(2-Ethoxyphenyl)purin-6,8-dion

2-(2-Butoxyphenyl)purin-6,8-dion

2-(2-Isobutoxyphenyl)purin-6,8-dion oder

15 2-(2-Allyloxyphenyl)purin-6,8-dion

oder ein pharmazeutisch verträgliches Salz davon bedeutet.

7. Verfahren gemäß Anspruch 1, wobei ein Formylierungsmittel aus Ameisensäure, C<sub>1-4</sub>-Alkylformiat, Formamid, C<sub>1-4</sub>-Alkylformamid, Formamidin, C<sub>1-4</sub>-Alkylformamidin oder Tri(C<sub>1-4</sub>)alkylorthoformiat ausgewählt wird.

20 8. Verfahren gemäß Anspruch 1, wobei ein Carbonylierungsmittel aus Harnstoff, Di(C<sub>1-4</sub>)alkylcarbonat, C<sub>1-4</sub>-Alkylchlorformiat, Phosgen, Trichlormethylchlorformiat oder Carbonyldiimidazol ausgewählt wird.

9. Verfahren zur Herstellung von 2-(2-Propoxyphenyl)-6-purinon oder eines pharmazeutisch verträglichen Salzes davon, umfassend die Umsetzung von 4,5-Diamino-2-(2-propoxyphenyl)pyrimidin-6-on oder einem Säureadditionssalz davon mit Formamid oder Formamidinacetat und gegebenenfalls die Bildung eines pharmazeutisch verträglichen Salzes davon.


25 10. Verfahren zur Herstellung von 2-(2-Propoxyphenyl)purin-6,8-dion oder eines pharmazeutisch verträglichen Salzes davon, umfassend die Umsetzung von 4,5-Diamino-2-(2-propoxyphenyl)pyrimidin-6-on mit Harnstoff oder Carbonyldiimidazol und gegebenenfalls die Bildung eines pharmazeutisch verträglichen Salzes davon.

30 11. Verfahren zur Herstellung eines Arzneimittels, umfassend die Vereinigung einer Verbindung der Formel (1), wie in Anspruch 1 definiert, oder eines pharmazeutisch verträglichen Salzes davon und eines pharmazeutisch verträglichen Trägers.

## 35 Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

40 1. composé de formule (1):



50 ou un de ses sels pharmaceutiquement acceptables, dans laquelle

R<sup>1</sup> est un groupe alkyle en C<sub>1-6</sub> ou alcényle en C<sub>2-6</sub>, et

R<sup>2</sup> est un atome d'hydrogène ou un groupe hydroxy.

2. Composé selon la revendication 1, dans lequel R<sup>2</sup> est un atome d'hydrogène.

3. Composé selon la revendication 1, dans lequel R<sup>2</sup> est un groupe hydroxy.

55 4. composé selon l'une quelconque des revendications 1 à 3, dans lequel R<sup>1</sup> est un groupe alkyle en C<sub>2-5</sub>.

5. Composé selon l'une quelconque des revendications 1 à 3, dans lequel R<sup>1</sup> est un groupe alcényle en

C<sub>3-6</sub>.

6. Composé selon la revendication 1 qui est :

la 2-(2-propoxyphényl)-6-purinone,

la 2-(2-éthoxyphényl)-6-purinone,

5 la 2-(2-butoxyphényl)-6-purinone,

la 2-(2-isobutoxyphényl)-6-purinone,

la 2-(2-propoxyphényl)purine-6,8-dione,

la 2-(2-méthoxyphényl)purine-6,8-dione,

la 2-(2-éthoxyphényl)purine-6,8-dione,

10 la 2-(2-butoxyphényl)purine-6,8-dione;

la 2-(2-isobutoxyphényl)purine-6,8-dione ou

la 2-(2-allyloxyphényl)purine-6,8-dione

ou un de ses sels pharmaceutiquement acceptables.

7. 2-(2-propoxyphényl)-6-purinone ou un de ses sels pharmaceutiquement acceptables.

15 8. 2-(2-propoxyphényl)purine-6,8-dione ou un de ses sels pharmaceutiquement acceptables.

9. Composé selon l'une quelconque des revendications 1 à 8 pour l'utilisation comme médicament.



10. Composé selon l'une quelconque des revendications 1 à 8 pour l'utilisation comme bronchodilatateur.

11. Composé selon l'une quelconque des revendications 1 à 8 pour l'utilisation comme vasodilatateur.

12. Composition pharmaceutique qui comprend un composé selon l'une quelconque des revendications


20 1 à 8 et un véhicule pharmaceutiquement acceptable.

13. Procédé pour préparer un composé de formule (1) ou un de ses sels pharmaceutiquement acceptables comme définis dans la revendication 1, qui comprend :

a) pour les composés dans lesquels R<sup>2</sup> est un atome d'hydrogène, la réaction d'un composé de formule (2) :35 dans laquelle R<sup>1</sup> est comme défini dans la revendication 1, avec un agent de formylation ;b) pour les composés dans lesquels R<sup>2</sup> est un groupe hydroxy, la réaction d'un composé de formule (2) comme précédemment défini avec un agent de carbonylation ;40 c) pour les composés dans lesquels R<sup>2</sup> est un atome d'hydrogène, la réaction d'un composé de formule (3) :dans laquelle R<sup>1</sup> est comme précédemment défini avec le 4-amino-5-imidazolecarboxamide,50 d) pour les composés dans lesquels R<sup>2</sup> est un atome d'hydrogène, la cyclisation d'un composé de formule (4) :

55

5




(4)

10 dans laquelle R¹ est comme précédemment défini ; puis, facultativement, la formation d'un sel pharmaceuti-  
quement acceptable.

14. Composé de formule (4) :

15



(4)


20

dans laquelle R¹ est un groupe alkyle en C₁-₈ ou alcényle en C₂-₆.

25 **Revendications pour les Etats contractants suivants : ES, GR**

1. Procédé pour la préparation d'un composé de formule (1) :

30




(1)

35

ou un de ses sels pharmaceutiquement acceptables, dans laquelle  
R¹ est un groupe alkyle en C₁-₈ ou alcényle en C₂-₆, et  
40 R² est un atome d'hydrogène ou un groupe hydroxy,  
lequel procédé comprend :

a) pour les composés dans lesquels R² est un atome d'hydrogène, la réaction d'un composé de formule (2) :

45



(2)


50

dans laquelle R¹ est comme précédemment défini, avec un agent de formylation ;

b) pour les composés dans lesquels R² est un groupe hydroxy, la réaction d'un composé de formule (2)  
comme précédemment défini avec un agent de carbonylation ;


55

c) pour les composés dans lesquels R² est un atome d'hydrogène, la réaction d'un composé de formule  
(3) :



(3)

dans laquelle R¹ est comme précédemment défini avec le 4-amino-5-imidazolecarboxamide,  
d) pour les composés dans lesquels R² est un atome d'hydrogène, la cyclisation d'un composé de formule (4) :



(4)

dans laquelle R¹ est comme précédemment défini ; puis, facultativement, la formation d'un sel pharmaceutiquement acceptable.

2. Procédé selon la revendication 1 pour préparer un composé dans lequel R² est un atome d'hydrogène.

3. Procédé selon la revendication 1 pour préparer un composé dans lequel R² est un groupe hydroxy.

4. Procédé selon l'une quelconque des revendications 1 à 3 pour préparer un composé dans lequel R¹ est un groupe alkyle en C<sub>2-5</sub>.

5. Procédé selon l'une quelconque des revendications 1 à 3 pour préparer un composé dans lequel R¹ est un groupe alcényle en C<sub>3-5</sub>.

6. Procédé selon la revendication 1 pour préparer un composé qui est :

la 2-(2-propoxyphényl)-6-purinone,

la 2-(2-éthoxyphényl)-6-purinone,

la 2-(2-butoxyphényl)-6-purinone,

la 2-(2-isobutoxyphényl)-6-purinone,

la 2-(2-propoxyphényl)purine-6,8-dione,

la 2-(2-méthoxyphényl)purine-6,8-dione,

la 2-(2-éthoxyphényl)purine-6,8-dione,

la 2-(2-butoxyphényl)purine-6,8-dione,

la 2-(2-isobutoxyphényl)purine-6,8-dione ou

la 2-(2-allyloxyphényl)purine-6,8-dione

ou un de ses sels pharmaceutiquement acceptables.

7. Procédé selon la revendication 1, dans lequel l'agent de formylation est choisi parmi l'acide formique, un formiate d'alkyle en C<sub>1-4</sub>, le formamide, un (alkyl en C<sub>1-4</sub>) formamide, la formamidine, une (alkyl en C<sub>1-4</sub>) formamidine ou un orthoformiate de tri(alkyle en C<sub>1-4</sub>).

8. Procédé selon la revendication 1, dans lequel l'agent de carbonylation est choisi parmi l'urée, un carbonate de di(alkyle en C<sub>1-4</sub>), un chloroformiate d'alkyle en C<sub>1-4</sub>, le phosgène, le chloroformiate de trichlorométhyle ou le carbonyldiimidazole.

9. Procédé pour préparer la 2-(2-propoxyphényl)-6-purinone ou un de ses sels pharmaceutiquement acceptables, qui comprend la réaction de la 4,5-diamino-2-(2-propoxyphényl)pyrimidine-6-one ou d'un de ses sels d'addition d'acides avec le formamide ou l'acétate de formamidine et facultativement la formation d'un sel pharmaceutiquement acceptable correspondant.

10. Procédé pour préparer la 2-(2-propoxyphényl)purine-6,8-dione ou un de ses sels pharmaceutiquement acceptables, qui comprend la réaction de la 4,5-diamino-2-(2-propoxyphényl)pyrimidine-6-one avec l'urée ou le carbonyldiimidazole et facultativement la formation d'un sel pharmaceutiquement acceptable correspondant.

11. Procédé pour préparer une composition pharmaceutique, qui comprend l'association d'un composé de formule (1) comme défini dans la revendication 1 ou d'un de ses sels pharmaceutiquement acceptables et d'un véhicule pharmaceutiquement acceptable.